PATENT . USSN 10/053,758 Docket 002980US: 018/183

P.003

## **CLAIM AMENDMENTS**

- (Currently amended) A monoclonal or recombinant antibody or <u>antigen binding</u> fragment thereof that <u>specifically</u> binds to human telomerase reverse transcriptase (hTRT) protein <u>having</u> the <u>sequence provided in SEQ. ID NO:225 (SEQ. ID NO:225)</u>.
- 2. (Currently amended) An antibody fragment that specifically binds to hTRT protein having the sequence provided in SEQ. ID NO:225 (SEQ. ID NO:225).
- 3. (Original) The antibody fragment of claim 2, which is an Fab fragment or an F(ab')2 fragment.
- 4. (Currently amended) The antibody or antigen binding fragment of claim 1, which is a chimeric human antibody.
- (Currently amended) The antibody or <u>antigen binding</u> fragment of claim 1, which has-a single chain is a single chain antibody.
- 6. (Currently amended) A pharmaceutical composition comprising the antibody or antigen binding fragment of claim 1 and a pharmaceutically acceptable carrier.
- 7. (Currently amended) The antibody or antigen binding fragment of claim 1, having a reporter molecule or label that is covalently or noncovalently bound.
- 8. (Currently amended) The antibody or <u>antigen binding</u> fragment of claim 7, wherein the reporter molecule or label is selected from an enzyme, a fluorescent agent, a chemiluminescent agent, a chromatogenic agent, and a magnetic particle.
- (Currently amended) A method of identifying a-polypeptide <u>hTRT</u> in a biological sample, comprising:
  - a) combining the biological sample with a monoclonal or recombinant antibody or <u>antigen</u> <u>binding</u> fragment thereof that <u>ean bind hTRT protein having the sequence provided in SEQ, ID NO:225</u> <u>specifically binds hTRT protein (SEQ, ID NO:225)</u>, under conditions where the antibody or fragment <u>will form forms</u> a complex with hTRT protein;
    - b) detecting complex formed as a result of a); and
  - c) identifying the sample as containing at-least-a pertion of hTRT protein if an antibody: protein complex is detected.

21-Sep-2004 07:25pm

P.004

T-393

- 10. (Original) The method of claim 9, which is an enzyme-linked immunosorbant assay method.
- 11. (Original) The method of claim 9, which is a radioimmunoassay method.
- 12. (Original) The method of claim 9, wherein the detecting comprises fluorescent activated cell sorting.
- 13. (Currently amended) A method of detecting an hTRT polypeptide in a blological sample, comprising:
  - a) combining the biological sample with a monoclonal or recombinant antibody or antigen binding fragment thereof according to claim 1, under conditions where an antibody will form forms a complex with hTRT protein (SEQ. ID NO:2); and
  - b) detecting any complex formed between the antibody or antigen binding fragment and hTRT protein the hTRT polypeptide.
- 14. (Original) The method of claim 13, which is an enzyme-linked immunosorbant assay method.
- 15. (Original) The method of claim 13, which is a radioimmunoassay method.
- 16. (Original) The method of claim 13, wherein the detecting comprises fluorescent activated cell sorting.
- 17. (Currently amended) A method of generating an antibody that specifically binds hTRT protein, comprising immunizing a host with a composition comprising a protein or poptide that contains an amino-acid-sequence selected from any 5-1100-centiguous-amino-acids in SEQ. ID NO:225 a composition comprising hTRT protein (SEQ. ID NO:225).

18 and 19. (Cancelled)

20. (Original) The method of claim 17, wherein the composition further comprises an adjuvant.

21 and 22. (Cancelled)